

## CONSTANT ACCELERATION FORMULAE



$$v = u + at$$

$$s = ut + 0.5at^2$$

$$s = vt - 0.5at^2$$

$$s = \frac{(u + v) \times t}{2}$$

$$v^2 = u^2 + 2as$$

v = final velocity (m/s)  
 u = initial velocity (m/s)  
 a = acceleration (m/s<sup>2</sup>)  
 s = displacement (m)  
 t = time (s)

## TORQUE



$$T = F \times d \times \sin(\theta)$$

T = Torque (Nm)  
 F = Force applied on lever (N)  
 d = Distance that the force is applied from the axis of rotation (m)  
 θ = Angle between force vector and distance vector, sin(θ) is equal to 1 if perpendicular

## NEWTON'S 2ND LAW OF MOTION



$$\Sigma F = m \times a$$

ΣF = net force acting on an object (N)  
 m = mass of object (kg)  
 a = acceleration of object (m/s<sup>2</sup>)


## MECHANICAL ENERGY



$$ME = U_s + U_g + KE$$

ME = mechanical energy (J)  
 U<sub>s</sub> = spring potential energy, a.k.a elastic potential energy (J)  
 U<sub>g</sub> = gravitational potential energy (J)  
 KE = kinetic energy (J)

## MOMENTUM



$$p = mv$$

p = Momentum (kg x m/s)  
 m = Mass (kg)  
 v = Velocity (m/s)

## GRAVITATIONAL POTENTIAL ENERGY



$$U_g = mgh$$

U<sub>g</sub> = gravitational potential energy (J)  
 m = Mass of object (kg)  
 g = Acceleration due to gravity = 9.8 m/s<sup>2</sup> = gravitational field strength = 9.8 N/kg  
 h = Vertical height above a reference level (m)

## TEMPERATURE CHANGE



$$Q = m \times C \times (T_{\text{final}} - T_{\text{initial}})$$

Q = heat energy applied (J)  
m = mass of object (kg)  
c = specific heat capacity of object (J/kg/K)  
T = initial and final temperatures of object (K)

## FORCE ON A MOVING CHARGE



$$F = qvb$$

F = Magnetic Force  
q = Charge of the particle  
v = Velocity of the charged particle  
b = Magnetic Field Strength


## OHM'S LAW



$$V = IR$$

V = Voltage (volts)  
I = Current (amps)  
R = Resistance (ohms)

## ELECTRICAL POWER



$$P = IV = I^2R = \frac{V^2}{R}$$

P = Power  
V = Voltage (volts)  
I = Current (amps)  
R = Resistance (ohms)

## KINETIC ENERGY



$$KE = \frac{1}{2} mv^2$$

KE = Kinetic energy of object (J)  
m = Mass of object (kg)  
v = Velocity of object (m/s)

## GRAVITATIONAL FORCE



$$F = \frac{G \times m_1 \times m_2}{r^2}$$

F = gravitational force of attraction from the center of mass of the two objects (N)  
G = gravitational constant  $6.674 \times 10^{-11}$  Nm kg  
m1 = mass of one of the objects (kg)  
m2 = mass of the other object (kg)  
r = distance between the center of the mass of the two objects (m)

## PRESSURE



$$P = \frac{F}{A}$$

P = Pressure (N/m or Pascals)  
F = Force applied on a surface (N)  
A = Area of surface (m<sup>2</sup>)

## ARCHIMEDES PRINCIPLE



$$F_b = F_{g \text{ of object}} = F_{g \text{ of fluid displaced}} = \rho_{\text{fluid}} \times V_{g \text{ of fluid displaced}} \times g$$

F<sub>b</sub> = Force of buoyance (N)  
F<sub>g of object</sub> = Weight of object (N)  
F<sub>g of fluid displaced</sub> = Weight of fluid displaced (N)  
ρ<sub>fluid</sub> = Density of fluid (kg/m<sup>3</sup>)  
V<sub>fluid displaced</sub> = Volume of fluid displaced by the object in the fluid (m<sup>3</sup>)  
g = gravitational field strength, a.k.a acceleration due to gravity = 9.8 N/kg = 9.8 m/s<sup>2</sup>

# Final Weeks to Enrol for Our January GAMSAT® Courses!

Enrolments Close Fri 10th Jan at 7 PM Melb Time



**Last chance to enrol for our January GAMSAT® Courses!**


**Enrol today and take advantage of our market-leading GAMSAT® Attendance Courses:**

- **Weekly Q&A Session with a Tutor** - Starting from the time of enrollment - every Wednesday night, our tutor will host an online Q&A session to discuss difficult Section 3 concepts, share perspectives on essay writing, and engage in live questions and answers with a group of students.
- **Extra Set of Virtual PBLs free of charge (for PBL students)**. For instance, if you enrol in January, you can choose to attend again in February. This offer applies to students who enrol for any Attendance PBLs. While the content remains consistent, revisiting the material will provide additional clarity and reinforcement for you.
- **75+ Contact Hours of live instruction** - Courses run by highly experienced tutors: Current medical students who have successfully completed the GAMSAT® Exam.
- **5 Realistic Mock Exam** - The mock-exams held at the end of the course are the most realistic in the industry and are a perfect rehearsal to the real GAMSAT® exam.
- **Industry-Leading Online Resources** - Get instant access to 5000+ MCQ Bank, 15 ACER-Like Online Exams that closely mimic the official ACER GAMSAT® exam, as well as performance tracking and percentile reporting comparing you to other GradReady students past and present.

***To Learn More or Enrol, Visit:***

***<https://gradready.com.au/gamsat-preparation-courses>***

## ELECTROSTATIC FORCE



$$F = \frac{k \times q_1 \times q_2}{r^2}$$

F = electrostatic force acting between two charged objects (N)

k = Coulomb's constant =  $9.0 \times 10 \text{ NmC}$

q1 = magnitude of one of the charged object (C)

q2 = magnitude of the other charged object (C)

r = distance between the center of the two charges (m)

## WORK



$$W = F \times d \times \cos(\theta)$$

W = Work done (J)

F = Force applied on object (N)

d = Distance object moves due to the application of the force (m)

$\theta$  = Angle between force vector and displacement vector,  $\cos(\theta)$  is equal to 1 when the force vector is in the same direction as the movement of the object

or

$$W = F \times d$$

W = Work done (J)

F = Component of the force that acts parallel to the distance only (N)

d = Distance object moves due to the application of the force (m)



Average Improvement of **20+** Percentile Points  
Over the last **10+** Years

**10,000+**  
Happy Students

**10+**  
Years Experience

Most Advanced  
Technology