

CONSTANT ACCELERATION FORMULAE



v = final velocity (m/s)
 u = initial velocity (m/s)
 a = acceleration (m/s²)
 s = displacement (m)
 t = time (s)

$$\begin{aligned}
 v &= u + at \\
 s &= ut + 0.5at^2 \\
 s &= vt - 0.5at^2 \\
 s &= \frac{(u + v) \times t}{2} \\
 v^2 &= u^2 + 2as
 \end{aligned}$$


NEWTON'S 2ND LAW OF MOTION



ΣF = net force acting on an object (N)
 m = mass of object (kg)
 a = acceleration of object (m/s²)

$$\Sigma F = m \times a$$

MOMENTUM



p = Momentum (kg x m/s)
 m = Mass (kg)
 v = Velocity (m/s)

$$p = mv$$

TORQUE



$$T = F \times d \times \sin(\theta)$$

T = Torque (Nm)
 F = Force applied on lever (N)
 d = Distance that the force is applied from the axis of rotation (m)
 θ = Angle between force vector and distance vector, $\sin(\theta)$ is equal to 1 if perpendicular

MECHANICAL ENERGY



$$ME = U_s + U_g + KE$$

ME = mechanical energy (J)
 U_s = spring potential energy, a.k.a elastic potential energy (J)
 U_g = gravitational potential energy (J)
 KE = kinetic energy (J)

GRAVITATIONAL POTENTIAL ENERGY



$$U_g = mgh$$

U_g = gravitational potential energy (J)
 m = Mass of object (kg)
 g = Acceleration due to gravity = 9.8 m/s² = gravitational field strength = 9.8 N/kg
 h = Vertical height above a reference level (m)

TEMPERATURE
CHANGE



$$Q = m \times C \times (T_{\text{final}} - T_{\text{initial}})$$

Q = heat energy applied (J)
m = mass of object (kg)
c = specific heat capacity of object (J/kg/K)
T = initial and final temperatures of object (K)

FORCE ON A MOVING
CHARGE



$$F = qvb$$

F = Magnetic Force
q = Charge of the particle
v = Velocity of the charged particle
b = Magnetic Field Strength


OHM'S
LAW



$$V = IR$$

V = Voltage (volts)
I = Current (amps)
R = Resistance (ohms)

ELECTRICAL
POWER



$$P = IV = I^2R = \frac{V^2}{R}$$

P = Power
V = Voltage (volts)
I = Current (amps)
R = Resistance (ohms)

KINETIC
ENERGY



$$KE = \frac{1}{2} mv^2$$

KE = Kinetic energy of object (J)
m = Mass of object (kg)
v = Velocity of object (m/s)

GRAVITATIONAL
FORCE



$$F = \frac{G \times m_1 \times m_2}{r^2}$$

F = gravitational force of attraction from the center of mass of the two objects (N)
G = gravitational constant 6.674×10^{-11} Nm kg
m1 = mass of one of the objects (kg)
m2 = mass of the other object (kg)
r = distance between the center of the mass of the two objects (m)

PRESSURE



$$P = \frac{F}{A}$$

P = Pressure (N/m or Pascals)
F = Force applied on a surface (N)
A = Area of surface (m²)

ARCHIMEDES
PRINCIPLE



$$F_b = F_{\text{g of object}} = F_{\text{g of fluid displaced}} = \rho_{\text{fluid}} \times V_{\text{g of fluid displaced}} \times g$$

F_b = Force of buoyancy (N)
F_{g of object} = Weight of object (N)
F_{g of fluid displaced} = Weight of fluid displaced (N)
ρ_{fluid} = Density of fluid (kg/m³)
V_{fluid displaced} = Volume of fluid displaced by the object in the fluid (m³)
g = gravitational field strength, a.k.a acceleration due to gravity = 9.8 N/kg = 9.8 m/s²

NOW YOU CAN JOIN AN EXTRA ROUND OF VIRTUAL PBLs FOR FREE!

Attend an Additional Round of PBL Classes at No Extra Cost!



At GradReady, we're constantly enhancing our offerings to better prepare you for the GAMSAT®. Now, when you enrol in any of our Attendance Courses, you can attend an additional round of Virtual PBLs of your choice at no extra charge.

For example, if you enrol in the August Course, you can attend again in February. Revisiting the material will provide you with extra clarity and reinforcement.


Our Virtual PBLs include:

- **15 Day Course, 75+ Contact Hours of Live Instruction** - Get support with Problem-Based Learning (PBL) Classes, Selective Topic Tutorials, and an Intensive Revision Weekend right before the exam.
- **Realistic Mock Exam** - The mock-exam held at the end of the course is the most realistic in the industry and is a perfect rehearsal to the real GAMSAT® exam. Our exam interface closely mimics the official ACER GAMSAT® exam, allowing you to familiarise yourself with the online delivery of the exam before test date!
- **Weekly Tutoring Session** - Every Monday night until the March GAMSAT®, our tutor will host an online tutoring session to discuss difficult reasoning concepts, share perspectives on essay writing, and engage in live questions and answers with a group of students.
- **Updated Class Curriculum** - Take advantage of our recently updated class curriculum with an increasing focus on reasoning skills, updated MCQ bank with new MCQs continuously being added, new content guides, and dedicated tutor Q&A time.
- **Learn from the Best** - Our courses are run by highly experienced tutors - current medical students who have successfully completed the GAMSAT® Exam.

Join 11,000+ GradReady Students who have improved their scores by 20+ Percentile Points on average, 11+ Years in a Row!

[Click Here](#) to Learn More & Sign Up Today. Spots Limited!

ELECTROSTATIC FORCE



$$F = \frac{k \times q_1 \times q_2}{r^2}$$

F = electrostatic force acting between two charged objects (N)

k = Coulomb's constant = $9.0 \times 10 \text{ NmC}$

q1 = magnitude of one of the charged object (C)

q2 = magnitude of the other charged object (C)

r = distance between the center of the two charges (m)

WORK



$$W = F \times d \times \cos(\theta)$$

W = Work done (J)

F = Force applied on object (N)

d = Distance object moves due to the application of the force (m)

θ = Angle between force vector and displacement vector, $\cos(\theta)$ is equal to 1 when the force vector is in the same direction as the movement of the object

or

$$W = F \times d$$

W = Work done (J)

F = Component of the force that acts parallel to the distance only (N)

d = Distance object moves due to the application of the force (m)



Average Improvement of **20+** Percentile Points
Over the last **10+** Years

10,000+
Happy Students

10+
Years Experience

Most Advanced
Technology

GRADREADY.COM.AU

TEL: (03) 9819 6696